

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Publications, etc. -- Nebraska Forest Service

Nebraska Forest Service

1999

Pennsylvania Urban and Community Forestry: Don't Top Trees

William Elmendorf

Penn State School of Forest Resources

Follow this and additional works at: <https://digitalcommons.unl.edu/nebforestpubs>



Part of the [Forest Sciences Commons](#)

Elmendorf, William, "Pennsylvania Urban and Community Forestry: Don't Top Trees" (1999). *Publications, etc. -- Nebraska Forest Service*. 71.

<https://digitalcommons.unl.edu/nebforestpubs/71>

This Article is brought to you for free and open access by the Nebraska Forest Service at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Publications, etc. -- Nebraska Forest Service by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.


Pennsylvania Urban and Community Forestry

Don't Top Trees



WHAT IS TOPPING?

- Ever notice trees in your community whose branches have been cut to stubs, causing them to look like sheared shrubs? These trees have been improperly topped. If you want healthy and safe trees, these examples of senseless destruction may provoke anger and disgust.



Tree butchers park the same way they prune—incorrectly!
USDA FOREST SERVICE

- Topping is the indiscriminate and drastic cutting of a tree branch or branches. It has been referred to as heading, stubbing, tipping, lopping, rounding-over, and dehorning. No matter what the name, topping disfigures trees and is detrimental to tree structure, health, and value.
- Trees are topped because they grow too large for the places where they were planted, or people become afraid of their large size. Trees are also topped because proper tree pruning methods are not used.

TOPPING HURTS A TREE IN MANY WAYS

- Death—In the most severe cases, topping can lead to the death of a tree.
- Unsafe structure—Topping causes the quick growth of dense, weak, upright branches called watersprouts. Sprouts caused by topping are often weakly attached. They grow so rapidly a tree can regain its original height in a short time with a dense and unwieldy crown.



Topping trees results in large areas of decay and fast-growing, weak, and unattractive watersprouts.

- Decline in health—Topping removes a tree's food production factory (leaves) and food stored in limbs that are cut. Topping also causes a tree to use valuable food stored in the trunk and roots to regrow limbs and branches. Topping redistributes the use of energy and has a substantial impact on the physiology of a tree. Although debated, topping also can

place a tree under stress and affect its tolerance to further injury.


- Ugliness—Topping replaces the natural form and beauty of a tree with unsightly branch stubs, large wounds, broom-like branch growth, and unhealthy branches.
- Value—Because of the negative impacts on appearance, structure, and health, topping can reduce the value of a large ornamental tree by thousands of dollars.
- Insects and diseases—The drastic removal of limbs by topping exposes remaining limbs to sunshine. This sudden exposure to light and heat can kill or damage a tree's living tissue (the bark and cambium) and is called sunscald. Trees have difficulty protecting themselves from sunscald and the large wounds that result from topping. These sunscald wounds are highly vulnerable to insect invasion and the spores and actions of decay fungi.

HOW TO PREVENT TOPPING

- Plant the right tree in the right place—Plant tree species that fit and will thrive in the places where they are planted. Don't plant large trees such as oaks, tulip poplar, and sycamore under power lines, in small tree lawns, in sidewalk cutouts, or close to buildings or signs.
- Buy quality nursery stock—To assure success when planting landscape trees, it is important to start with a healthy plant that has good structural form. Characteristics that are important to consider when purchasing nursery stock can be found in the American Standards for Nursery Stock, available from the American Association of Nurserymen, (202) 789-2900.

■ Prune young trees—*The most important time to prune trees is in the first 5 to 10 years of their life.* Proper pruning of young trees promotes correct development of structure, saves money by removing branches when they are small, and keeps pruning wounds small.

■ Prune correctly—All tree pruning should be performed in accordance with current International Society of



Proper pruning enhances and protects the natural shape of a tree.
International Society of Arboriculture

Arboriculture tree maintenance standards. Insist that thinning cuts and natural target pruning techniques are used.


■ Proper utility clearance—Because of the vigorous growth of sprouts, topping makes utility line clearance more difficult, frequent, and expensive. Most utilities use an arboriculturally correct method called directional pruning instead of topping. Directional pruning directs subsequent growth away from wires and leaves no branch stubs.

■ Hire a qualified arborist—Although tree topping may cost less and take less time, why pay for an inferior service? Hire a qualified arborist, preferably a certified arborist, who knows about proper and safe tree care techniques. Do not blindly accept the lowest price. In tree work, the

old adage 'you get what you pay for' is usually true. Ask for customer references and proof of liability insurance and worker's compensation insurance.

■ Gather good information—For more information on tree care and management, contact the Penn State Cooperative Extension Urban Forestry Program at (814) 863-7941, the Pennsylvania Urban and Community Forestry Council at (717) 783-0385, the International Society of Arboriculture at (217) 355-9411, the National Arborist Association at (603) 673-3311, the National Arbor Day Foundation, at (402) 474-5655, or your local electric utility arborist.

Thanks to the International Society of Arboriculture and the National Arbor Day Foundation for information and illustrations used in this fact sheet.



When required because of obstructions, a tree's height should be reduced using thinning cuts.
International Society of Arboriculture

Produced by William Elmendorf, Instructor of Community Forestry and Henry Gerhold, Professor of Forest Genetics, Penn State School of Forest Resources.

This fact sheet was prepared by Penn State with guidance from the Pennsylvania Urban and Community Forestry Council and support from the Pennsylvania DCNR Bureau of Forestry. For more information, contact the Pennsylvania Urban and Community Forestry Program, School of Forest Resources, The Pennsylvania State University, 108 Ferguson, University Park, PA 16802; (814) 863-7941.

This publication is available in alternative media on request.

The Pennsylvania State University is committed to the policy that all persons shall have equal access to programs, facilities, admission, and employment without regard to personal characteristics not related to ability, performance, or qualifications as determined by University policy or by state or federal authorities. The Pennsylvania State University does not discriminate against any person because of age, ancestry, color, disability or handicap, national origin, race religious creed, sex, sexual orientation, or veteran status. Direct all inquiries regarding the nondiscrimination policy to the Affirmative Action Director, The Pennsylvania State University, 201 Willard Building, University Park PA 16802-2801; Tel. (814) 863-4700/V; (814) 863-1150/TTY.



Pennsylvania
Urban and Community
Forestry Council

